Variation in the nutrient composition of different feedstuffs for horses

Frieder J. Schwarz¹, Heike Sliwinski¹, Manfred Schuster² and Ewald Rosenberger²

Department of Animal Science, Division of Animal Nutrition, Technical University of Munich, Freising-Weihenstephan¹ and Bavarian State Research Center for Agriculture, Institute of Animal Breeding, Poing²

Introduction

Calculating rations for horses requires the most accurate knowledge possible of the energy and nutrient contents of the individual feedstuffs. In general, average figures from tables of nutritional values (in Germany: DLG-Futterwerttabelle für Pferde, 1995) are used. However, large deviations can be found, leading to possible under- or over-supply of energy or nutrients despite ration planning. This wide ranging investigation of the most common feedstuffs used for horses should provide a general picture of energy and nutrient contents in rations under practical farm conditions.

Material and methods

Three samples of feedstuffs from forty farms/stables, evenly distributed over Bavaria, were taken, at 4-week intervals during 2001/2002. These individual samples were combined for each of the feedstuff of each stable. The separate ration components investigated were: forage (meadow hay, 1st and 2nd cut, lucerne hay), silages (grass silage, 1st, 2nd and subsequent cuts, ryegrass-silage, cereal whole-crop silage, maize silage), grass cobs and maize cobs, cereals (oats, barley, maize) and proprietary horse food/mixes. A total of 159 feedstuffs were analysed, the crude nutrients by the Weende analysis chemically and using the NIRS system. Furtheron, the samples were wet-ashed with acid and then analysed for major and trace elements by AAS. Phosphorus content was determined by colour reaction with an ammoniavanadate/heptamolybdate solution. Digestible crude protein and digestible energy contents were calculated using the analysed crude nutrients and digestibility coefficients for horses from the DLG-Futterwerttabelle (1995).

Results

In order to get a better general view of the analysed values, only the averages for the conserved grass products (meadow hay, 1st and 2nd cut, grass silage, 1st, 2nd and subsequent

cuts, grass cobs) and for the cereals (oats, barley, maize) are given, together with the minimum and maximum values showing the range. Table 1 shows the average contents of dry matter, crude nutrients and energy, table 2 the major element contents and table 3 the trace element values.

Table 1 Mean dry matter (DM), crude nutrients (crude ash (CA), crude protein (CP), crude fibre (CF)), digestible crude protein (DP) and digestible energy (DE) of different feedstuffs.

	DM (g/kg)	CA (g/kg DM)	CP (g/kg DM)	CF (g/kg DM)	DP (g/kg DM)	DE (MJ/kg DM)
Hay, 1 st cut (n = 43)	848 795-24876	60 40-76	78 58-121	331 294-361	46 31-76	8.96 7.6-10.1
Hay, 2 nd cut (n = 3)	826 824-830	68 43-83	104 97-106	295 293-297	64 60-67	9.85 9.7-10.2
Grass silage 1 st cut (n = 12)	571 420-755	79 53-129	104 62-160	322 268-372	71 42-112	10.0 9.5-11.2
Grass silage 2 nd cut (n = 24)	540 285-769	130 92-218	123 92-184	259 204-294	89 61-138	9.45 8.2-11.1
Grass cobs (n = 7)	906 888-924	127 89-247	114 95-160	241 171-210	97 64-138	9.86 12.5-13.4
Oats (n = 34)	870 851-886	29 25-31	115 95-136	116 96-152	91 75-107	13.0 12.5-13.4
Barley (n = 10)	873 859-596	23 22-25	117 98-126	51 49-56	86 72-91	14.5 14.5-14.6
Maize	877 861-887	16 14-18	98 90-108	22 19-25	68 61-75	15.6 15.5-15.6

Table 2 Mean content of major minerals of different feedstuffs.

	Ca (g/kg DM)	P (g/kg DM)	Mg (g/kg DM)	Na (g/kg DM)	K (g/kg DM)
Hay, 1 st cut (n = 43)	4.15 1.9-7.2	2.25 1.2-3.6	1.63 1.0-2.3	0.38 0.03-2.0	16.9 7.8-33.0
Hay, 2 nd cut (n = 3)	5.89 5.0-7.1	3.10 2.9-3.5	2.07 1.6-2.6	0.48 0.05-0.8	20.5 10.9-25.7
Grass silage 1 st cut (n = 12)	5.48 2.4-11.6	3.06 2.2-5.0	1.91 1.2-3.9	0.20 0.05-0.6	25.7 16.1-40.0
Grass silage 2 nd cut (n = 24)	7.86 3.5-16.9	3.44 2.2-4.9	2.75 1.6-4.8	0.29 0.03-2.4	20.9 13.5-33.4
Grass cobs (n = 7)	8.01 5.6-11.0	3.22 2.5-3.7	2.85 1.9-3.3	0.50 0.29-1.0	21.0 17.6-24.9
Oats (n = 34)	0.88 0.61-1.6	3.79 3.4-4.1	1.20 1.1-1.5	0.07 0.02-0.3	4.93 4.0-6.8
Barley (n = 10)	0.53 0.39-0.69	3.94 3.6-4.5	1.23 1.1-1.3	0.07 0.04-0.1	5.64 4.5-6.1
Maize (n = 9)	0.20 0.06-0.64	3.25 2.9-3.5	1.27 1.1-1.4	0.07 0.02-0.2	3.68 1.4-4.2

Table 3 Mean content of trace elements of different feedstuffs.

	Cu (mg/kg)	Mn (mg/kg DM)	Zn (mg/kg DM)	Se (µg/kg DM)
Hay, 1 st cut (n = 43)	4.60 3.2-6.0	97.0 27.3-331	30.9 17.5-44.0	23.6 8.2-91.0
Hay, 2 nd cut (n = 3)	5.41 5.0-5.8	47.5 31.7-64.7	36.4 16.9-47.5	21.0 12.5-36.1
Grass silage 1 st cut (n = 12)	5.50 3.5-7.7	110 36.9-201	31.4 16.9-47.5	19.1 9.3-48.6
Grass silage 2 nd cut (n = 24)	7.18 6.0-10.4	169 28.6-410	40.9 27.5-73.1	48.4 10.4-197
Grass cobs (n = 7)	7.89 6.7-10.0	185 100-306	42.3 34.9-54.2	46.1 21.6-81.3
Oats (n = 34)	3.89 2.5-5.1	43.5 17.7-73.2	23.8 15.8-37.6	15.9 6.4-33.4
Barley (n = 10)	4.42 2.2-5.5	17.6 11.8-24.5	25.8 20.6-33.2	14.4 9.6-21.5
Maize (n = 9)	2.58 1.9-3.9	6.82 5.2-10.8	20.7 17.1-25.7	13.8 4.4-40.2

Discussion

Hay is one of the main constituents in any ration for horses. 43 samples of 1st cut meadow hay were analysed, and only three of 2nd cut, showing that hardly any 2nd cut hay is used.

The relatively high contents of crude fibre (33% in the DM), coupled with the very low values for crude protein (8% in the

DM) suggest a late harvest date. As rising crude fibre content leads to a reduction in digestibility, the digestible energy value of just under 9.0 MJ/kg DM is rather low. The average contents of major minerals and trace elements, with the exception of manganese, are also on the low side, and the very large range must be mentioned here.

Grass silage (n=36) is now an important component in rations for horses in many stables, with 2nd and further cuts often used for silage. However, the DM content of the silages varied enormously between below 30% and more than 70%, so that we cannot yet say that there are set methods for wilting and ensiling silages for horses. Also, a distinction must be made between silages (DM 45-55%) with normal fermentation and products with higher DM contents (haylage, DM >65%). Both products require high compression for successful conservation, so that forages in an advanced physiological state are unsuitable. In general, grass silages have higher energy and mineral contents than hay.

Grass cobs are produced by drying the grass using fossil fuels, with subsequent compression into cobs. This process is expensive, requiring a high quality product to justify the

expense, but in some cases a high content of ash (25%) was found, which reduces the nutritional value significantly.

The separate samples of barley and maize varied only little in contents of crude nutrients, energy and minerals. There were, however, big variations in the crude protein and crude fibre contents of oats, therefore leading to a wide range of energy contents. All cereal samples showed low major and trace element contents relative to the requirement of horses.

References

DLG-Futterwerttabelle für Pferde (1995): Universität Hohenheim, Dokumentationsstelle. 3rd ed., DLG-Verlag, Frankfurt

F. J. Schwarz
Department of Animal Science
Division of Animal Nutrition
Technical University of Munich
85350 Freising-Weihenstephan
Germany
schwarzf@wzw.tum.de